arXiv:1209.4399v1 [cond-mat.str-el] 20 Sep 2012

Three dimensional Symmetry Protected Topological Phase close to Antiferromagnetic Néel order

Cenke Xu[1]

[1]*Department of Physics, University of California, Santa Barbara, CA 93106*

(Dated: September 21, 2012)

It is well-known that the Haldane phase of one-dimensional spin-1 chain is a symmetry protected topological (SPT) phase, which is described by a nonlinear Sigma model (NLSM) with a $\Theta-$term at $\Theta = 2\pi$. In this work we study a 3+1 dimensional SPT phase of SU($2N$) antiferromagnetic spin system with a self-conjugate representation on every site. The spin ordered Néel phase of this system has a ground state manifold $\mathcal{M} = \frac{\mathrm{U}(2N)}{\mathrm{U}(N)\times\mathrm{U}(N)}$, and this system is described by a NLSM defined with manifold $\mathcal{M}$. Since the homotopy group $\pi_4[\mathcal{M}] = \mathbb{Z}$ for $N > 1$, this NLSM can naturally have a $\Theta-$term. We will argue that when $\Theta = 2\pi$ this NLSM describes a SPT phase. We will also construct a 3d lattice model for this SPT phase, whose long wavelength physics is precisely described by this NLSM with $\Theta = 2\pi$. This SPT phase is protected by the SU($2N$) spin symmetry, or its subgroup SU(N)$\times$SU(N) $\rtimes Z_2$, without assuming any lattice symmetry.

PACS numbers:

1. INTRODUCTION

In quantum many-body systems, quantum mechanics can lead to at least *three* types of exotic/nontrivial quantum disordered phases: (1) Topological phases with a gapped spectrum and bulk topological degeneracy, (2) algebraic liquid phases with gapless bulk spectrum and power-law correlations, and (3) symmetry protected topological phases.

A symmetry protected topological (SPT) phase is a state of matter with gapped and nondegenerate bulk spectrum, but cannot continuously evolve into a direct product state without a bulk phase transition, when and only when the Hamiltonian of the entire evolution is invariant under certain global symmetry G [1]. In terms of its phenomena, a SPT phase on a $d-$dimensional lattice should satisfy at least the following three criteria:

(1). On a $d-$dimensional lattice without boundary, this phase is fully gapped, and nondegenerate;

(2). On a $d-$dimensional lattice with a $(d-1)-$dimensional boundary, if the Hamiltonian of the entire system (including both bulk and boundary Hamiltonian) preserves certain symmetry G, this phase is either gapless, or gapped but degenerate.

(3). The boundary state of this $d-$dim system cannot be realized as a $(d-1)$-dim lattice system built with the same onsite Hilbert space, and with the same symmetry G.

If a $d-$dim SPT phase satisfies all three criteria (1), (2) and (3), this phase is a SPT. Both the $2d$ quantum spin Hall insulator [2–4] and $3d$ Topological insulator [5–7] are perfect examples of SPT phases protected by time-reversal symmetry.

Notice that the second criterion (2) implies the following two possibilities: On a lattice with a boundary, the system is either ($2a$) gapless, or ($2b$) gapped but degenerate. When $d \geq 3$, the degeneracy of ($2b$) can correspond to either spontaneous breaking of G, or correspond to certain topological degeneracy at the boundary. Which case occurs in the system will depend on the detailed Hamiltonian at the boundary of the system. For example, with interaction, the edge states of $2d$ QSH insulator, and $3d$ TBI can both be gapped out through spontaneous symmetry breaking at the boundary, and this spontaneous symmetry breaking can occur through a boundary transition, without destroying the bulk state [8–10].

In this work we will focus on *bosonic* spin systems. The simplest example of SPT phase of spin system is the Haldane phase of one dimensional spin-1 chain. Unlike the free fermion case, although there is a classification of bosonic SPT using group cohomology [1], specific models of higher dimensional bosonic spin systems are not well understood. So far, in most studies, construction of 2d and 3d bosonic SPTs has been focused in systems with U(1) symmetry [11–13]. In this work we will study a 3+1 dimensional analogue of the Haldane phase, which is constructed as a SU($2N$) spin state with a self-conjugate representation on each site. Just like the Haldane phase, this 3+1d SPT is described by a nonlinear sigma model defined with semiclassical antiferromagnetic order parameter plus a topological $\Theta-$term.

2. HALDANE PHASE

Although symmetry protected topological phase is a pure quantum phenomenon without any classical analogue, the Haldane phase can still be described semiclassically by a nonlinear Sigma model (NLSM), which is defined *only* in terms of the semiclassical Néel order parameter $\vec{n}$ [14, 15]:

$$\mathcal{L} = \frac{1}{g}(\partial_\mu \vec{n})^2 + \frac{i\Theta}{8\pi}\epsilon_{abc}\epsilon_{\mu\nu} n^a \partial_\mu n^b \partial_\nu n^c. \qquad (1)$$

When the system has SO(3) symmetry, the entire manifold of the configurations of Néel order parameter is S^2. If we assume the trivial vacuum has $\Theta = 0$, then when $\Theta = 2\pi$, this model describes the Haldane phase. At the boundary of the $1d$ system, *i.e.* the domain wall between $\Theta = 0$ and 2π, the $\Theta-$term reduces to a 0+1d Wess-Zumino-Witten (WZW) term at level 1, whose ground state has two fold degeneracy [16]. As long as the SO(3) symmetry is preserved, this two fold degeneracy at the boundary cannot be lifted. Thus by definition the Haldane phase is a SPT protected by SO(3) spin rotation symmetry.

Now let us couple two Haldane phases to each other:

$$\mathcal{L} = \frac{1}{g}(\partial_\mu \vec{n}_1)^2 + \frac{i2\pi}{8\pi}\epsilon_{abc}\epsilon_{\mu\nu} n_1^a \partial_\mu n_1^b \partial_\nu n_1^c + 1 \rightarrow 2 + A(\vec{n}_1 \cdot \vec{n}_2). \quad (2)$$

When $A = -\infty$, effectively $\vec{n}_1 = \vec{n}_2 = \vec{n}$, then the system is effectively described by an O(3) NLSM with $\Theta = 4\pi$; while when $A = +\infty$, the effective NLSM for the system has $\Theta = 0$. When parameter A is tuned from $-\infty$ to $+\infty$, the entire phase diagram with $A \in (-\infty, +\infty)$ is gapped in the bulk. Thus the theory with $\Theta = 4\pi$ and $\Theta = 0$ are equivalent. This analysis implies that with SO(3) symmetry, $1d$ spin systems have two different classes: there is a trivial class with $\Theta = 4\pi k$, and a nontrivial Haldane class with $\Theta = (4k+2)\pi$. This Z_2 classification is consistent with the cohomology formalism developed in Ref. [1]. In fact, using the renormalization group calculation [17–20], and the general nonperturbative argument in Ref. [21], we can derive a phase diagram for model Eq. 1: The system is topological when $\Theta \in ((4k+1)\pi, (4k+3)\pi)$, while the system is trivial when $\Theta \in ((4k-1)\pi, (4k+1)\pi)$; $\Theta = (2k+1)\pi$ is the transition, where the system is either gapless, or two fold degenerate.

There are various ways of describing the Haldane phase on the lattice. In the follows we will choose one particular description that will be generalized to higher dimensions later. The Haldane phase can be described on the lattice as follows: On every site we introduce a slave fermion with both spin-1/2 index and SU(2) color index: $f_{i,A,\alpha}$, and the spin-1 operator is represented as [22]

$$\vec{S}_j = \frac{1}{2}\sum_{A=1}^{2} f^\dagger_{j,A,\alpha}\vec{\sigma}_{\alpha\beta} f_{j,A,\beta}. \quad (3)$$

In order to match the slave fermion Hilbert space with the spin-1 Hilbert space, we have to impose two different constraints on each site:

$$\sum_{\alpha,A} f^\dagger_{i,A,\alpha} f_{i,A,\alpha} = 2, \quad \sum_{\alpha,A,B} f^\dagger_{j,A,\alpha}\rho^\mu_{AB} f_{j,B,\alpha} = 0, \quad (4)$$

where ρ^μ are three Pauli matrices of the color space. The second constraint guarantees that on every site the color space is in a total antisymmetric singlet, thus the spin is in a total symmetric spin-1 representation. Then the Haldane phase corresponds to the following mean field state of $f_{A,\alpha}$: $f_{1,\alpha}$ forms valence bonds on links $(2i, 2i+1)$, while $f_{2,\alpha}$ forms valence bonds on links $(2i+1, 2i+2)$ (Fig. 1a). In terms of low energy field theory of the slave fermion, the Haldane phase is described by the following Lagrangian:

$$\mathcal{L} = \bar{\psi}\gamma_\mu\partial_\mu\psi + m_4\bar{\psi}\rho^z\psi. \quad (5)$$

Here the Dirac fermion ψ is the low energy mode of f, which is expanded at the two Fermi points $k_f = \pm\pi/2$ in the 1d Brillouin zone. If we couple the Néel order parameter to the slave fermion,

$$(-1)^j \vec{n}_j \cdot \sum_A f^\dagger_{j,A,\alpha}\vec{\sigma}_{\alpha\beta} f_{j,A,\beta} \sim \vec{n}\cdot\bar{\psi}\gamma_5\vec{\sigma}\psi, \quad (6)$$

Eq. 1 can be readily derived after integrating out the slave fermions [23]. Notice that in Eq. 5 the gauge fields introduced by constraints Eq. 4 are ignored, but in 1+1 dimension gauge fields are always confining.

The Haldane phase is a SPT *only* when the color-singlet constraint $f^\dagger_j \rho^\mu f_j = 0$ is strictly imposed on every site, *i.e.* when the Hilbert space on every site is rigorously spin-1. If this constraint is given up, the Hilbert space on every site is enlarged to 6 dimension, and the Haldane phase becomes trivial, because it can now be adiabatically connected to a direct product state with spin-singlet on every site. Actually, besides the Haldane phase mass gap m_4, we can consider another mass gap $m_5\bar{\psi}\gamma_5\rho^z\psi$ of the Dirac fermion ψ. Physically m_5 corresponds to a "color density wave" on the lattice, which is not allowed if the color singlet constraint is imposed strictly on every site. Without the color singlet constraint, the Haldane phase can adiabatically evolve into the color density wave state, by turning on m_5.

Eq. 1 already assumed a rigorous spin-1 Hilbert space, since it does not involve any orbital degree of freedom besides vector $\vec{n}$.

3. 3D SPT OF SU($2N$) SPIN SYSTEM

3.1 Field Theory Description

Let us try to look for higher dimensional generalizations of the Haldane phase of spin-1 chain, which has a description in terms of NLSM plus a $\Theta-$term. The most naive generalization would be the AKLT state in higher dimensions, for instance the spin-2 AKLT phase on the square lattice. The boundary of the spin-2 AKLT phase on the square lattice is a spin-1/2 chain, which according to the LSM theorem cannot be gapped and nondegenrate, thus the spin-2 AKLT state seems to be a SPT. However, in order to protect the spin-2 AKLT state, we need translation symmetry, since the boundary spin-1/2 chain can



FIG. 1: (a). The pictorial representation of the Haldane phase, where the blue and red links stand for the valence bonds of slave fermions with color $A = 1$ and 2 respectively. (b) A domain wall of Θ' on the boundary of our $3d$ SPT. (c) The Young diagram of the self-conjugate representation of the SU($2N$) spin system that we are considering.

be dimerized and gapped out once the translation symmetry of the system is explicitly broken. Thus this is not an ideal generalization of the $1d$ Haldane phase, whose stability does not rely on any translation symmetry. The spin-2 AKLT state on the square lattice can indeed be described by a NLSM with a topological term, but the configurational space of this NLSM would involve both the Néel and dimerization order parameters.

The goal of this paper is to find a three dimensional SPT phase *without* assuming the translation symmetry. Inspired by Eq. 1, we should first look for magnetic systems, whose ground state manifold $\mathcal{M}$ of the spin ordered phase has a nontrivial homotopy group $\pi_4[\mathcal{M}] = \mathbb{Z}$, then this SPT can be described by a NLSM defined in manifold $\mathcal{M}$ with a Θ term. The SU($2N$) antiferromagnet with self-conjugate representation satisfies this criterion: its magnetic ordered phase has GSM [24, 25]

$$\mathcal{M} = \frac{\mathrm{U}(2N)}{\mathrm{U}(N) \times \mathrm{U}(N)}, \quad \pi_4[\mathcal{M}] = \mathbb{Z}, \text{ for } N \geq 2. \qquad (7)$$

Every Néel order configuration $\mathcal{P} \in \mathcal{M}$ can be represented as

$$\mathcal{P} = V^\dagger \Omega V, \quad \Omega = \begin{pmatrix} \mathbf{1}_{N\times N}, & \mathbf{0}_{N\times N} \\ \mathbf{0}_{N\times N}, & -\mathbf{1}_{N\times N} \end{pmatrix} \qquad (8)$$

where V is a SU($2N$) matrix. $\mathcal{P}$ is a hermitian traceless order parameter that satisfies $\mathcal{P}^2 = \mathbf{1}$. In fact, when $N = 1$, $\mathcal{M}$ is precisely S^2, and $\mathcal{P}$ can always be represented as $\mathcal{P} = \vec{n} \cdot \vec{\sigma}$, where $\vec{n}$ is the Néel vector.

Since $\pi_4[\mathcal{M}] = \mathbb{Z}$, the following NLSM defined on $\mathcal{M}$ can be written down:

$$\mathcal{S} = \int d^3x d\tau \ \frac{1}{g} \mathrm{tr}[\partial_\mu \mathcal{P} \partial_\mu \mathcal{P}] + \frac{i\Theta}{256\pi^2} \mathrm{tr}[\mathcal{P}\partial_\mu \mathcal{P} \partial_\nu \mathcal{P} \partial_\rho \mathcal{P} \partial_\lambda \mathcal{P}] \epsilon_{\mu\nu\rho\lambda}. \qquad (9)$$

By tuning the parameter g, there is obviously an order-disorder transition. When g is small, the system is in a spin ordered phase where $\mathcal{P}$ is condensed and spontaneously breaks the SU($2N$) symmetry; when g is large, the system is in a disordered phase, and this disordered phase is what we are interested in.

In the follows we will focus on the disordered phase of Eq. 9 with $\Theta = 2\pi$, while assuming the trivial vacuum of this spin system has $\Theta = 0$. With $\Theta = 2\pi$, the bulk spectrum of the field theory is identical to $\Theta = 0$, thus the disordered phase is gapped and nondegenerate. Then we can safely integrate out the bulk, and look at the boundary theory. Using the standard bulk-boundary correspondence, we can derive the boundary theory of Eq. 9, which is a 2+1 dimensional NLSM defined in $\mathcal{M}$ with a WZW term at level $k = 1$:

$$\mathcal{S}_b = \int d^2x d\tau \ \frac{1}{g} \mathrm{tr}[\partial_\mu \mathcal{P} \partial_\mu \mathcal{P}] + \int_0^1 du \int d^2x d\tau \ \frac{i2\pi k}{256\pi^2} \mathrm{tr}[\mathcal{P}\partial_\mu \mathcal{P} \partial_\nu \mathcal{P} \partial_\rho \mathcal{P} \partial_\lambda \mathcal{P}] \epsilon_{\mu\nu\rho\lambda} \qquad (10)$$

Here $u \in (0, 1)$, and $\mathcal{P}(\vec{x}, \tau, u)$ is an extension of $\mathcal{P}(\vec{x}, \tau)$ that satisfies

$$\mathcal{P}(\vec{x}, \tau, 1) = \mathcal{P}(\vec{x}, \tau), \quad \mathcal{P}(\vec{x}, \tau, 0) = \Omega. \qquad (11)$$

Such WZW terms can be analyzed very reliably in 0+1d and 1+1d, and in both cases, these terms change the ground state dramatically. In 0+1d, a WZW term leads to degenerate ground states; in 1+1d, it drives the system to a stable gapless fixed point described by conformal field theory [26, 27]. In higher dimensions, nontrivial effects of a WZW term are still expected, but we no longer have a complete understanding. In the follows I will argue that the disordered phase of the 2+1d boundary theory Eq. 10 must be either gapless or degenerate.

In order to make this argument, let us first weakly break the SU($2N$) symmetry down to SU(N)$\times$SU(N)$\rtimes Z_2$. This residual SU(N)$\times$SU(N) symmetry transformation can be written as

$$V = \begin{pmatrix} V_L, & \mathbf{0}_{N\times N} \\ \mathbf{0}_{N\times N}, & V_R \end{pmatrix} \qquad (12)$$

while the residual Z_2 symmetry corresponds to exchanging V_L and V_R. With this symmetry reduction, the order parameter $\mathcal{P}$ can be written as

$$\mathcal{P} = \begin{pmatrix} \cos(\theta)\mathbf{1}_{N\times N}, & i\sin(\theta) U \\ -i\sin(\theta) U^\dagger, & -\cos(\theta)\mathbf{1}_{N\times N} \end{pmatrix}, \qquad (13)$$

where U is an SU(N) matrix. Under the transformation V_L and V_R, U transforms $U \to V_L^\dagger U V_R$, thus the

SU(N)×SU(N) residual symmetry precisely corresponds to the left and right transformation of U. Under the Z_2 symmetry transformation,

$$Z_2: \quad \theta \to \pi - \theta, \quad U \to U^\dagger. \tag{14}$$

With this symmetry reduction, the boundary theory with the WZW term Eq. 10 is reduced to a principal chiral model (PCM) defined on manifold SU(N) with a Θ' term:

$$\begin{aligned} \mathcal{S}_b \ \to \ & \int d^2x d\tau \ \frac{1}{g} \mathrm{tr}[\partial_\mu U^\dagger \partial_\mu U] \\ + \ & \frac{i\Theta'}{24\pi^2} \mathrm{tr}[U^\dagger \partial_\mu U U^\dagger \partial_\nu U U^\dagger \partial_\rho U]. \end{aligned} \tag{15}$$

If the Z_2 symmetry of the residual symmetry is unbroken, *i.e.* the expectation value of θ is $\pi/2$, then the derived boundary SU(N) PCM has precisely $\Theta' = \pi$. When this Z_2 symmetry is explicitly broken, namely at the boundary we turn on a background field that tunes θ away from $\pi/2$, then a straightforward calculation shows that the derived Θ' is also tuned away from π, and $\Theta' = 2\pi(2 + \cos(\theta))(\sin(\theta/2))^4$. Under the Z_2 transformation, $\Theta' \to 2\pi - \Theta'$.

The phase diagram of 2+1d SU(N) PCM was studied in Ref. [21], where it was argued that when g is large enough, $\Theta' > \pi$ and $\Theta' < \pi$ are two different disordered phases, which are both fully gapped and nondegenerate. These two disordered phases are separated by either a first or second order transition at $\Theta' = \pi$. Thus at $\Theta' = \pi$ the system must be either gapless or two fold degenerate. In our formalism we can see that the residual SU(N)×SU(N)⋊Z_2 symmetry guarantees $\Theta' = \pi$ at the boundary, *i.e.* these symmetries guarantee the boundary of Eq. 9 cannot be trivially gapped out.

Just like the 1d case discussed before, the existence of a Θ'−term in a 2+1d PCM can lead to nontrivial edge states. Since the 2d boundary of a 3d system cannot have its own boundary, the edge states can only exist at a domain wall of physical order parameters. Now let us create the following configuration of Θ' at the 2d boundary (Fig. 1b):

$$\begin{aligned} \Theta'(\vec{x}) \ &= \ 2\pi, \quad \text{for } |\vec{x}| < R, \\ \Theta'(\vec{x}) \ &= \ 0, \quad \text{for } |\vec{x}| > R. \end{aligned} \tag{16}$$

According to our previous work [21], in the disordered phase of Eq. 15, at the domain wall $|\vec{x}| = R$, there is a gapless SU$(N)_L$ × SU$(N)_R$ conformal field theory with level $k = 1$. The SU$(N)_L$ and SU$(N)_R$ charges move clockwise and counter-clockwise respectively along the domain wall. Later these domain wall states will help to argue that with the SU(N)×SU(N)⋊Z_2 symmetry, model Eq. 15 can only be realized at the boundary of a 3d system, *i.e.* Eq. 9 describes a 3d SPT.

By coupling two copies of Eq. 9 together like Eq. 2, one can show that the theory with $\Theta = 4\pi$ can be continuously connected to $\Theta = 0$ without a bulk transition. Thus again Eq. 9 describes a 3d SPT with Z_2 classification: there is a trivial class with $\Theta = 4\pi k$, and a nontrivial Haldane class with $\Theta = (4k + 2)\pi$. Using the same argument as Ref. [21], we can derive a phase diagram for model Eq. 9: The system is topological when $\Theta \in ((4k + 1)\pi, (4k + 3)\pi)$, while the system is trivial when $\Theta \in ((4k - 1)\pi, (4k + 1)\pi)$; $\Theta = (2k + 1)\pi$ is the transition, where the system is either gapless, or two fold degenerate.

3.2 Lattice Construction

Now let us construct a spin state for Eq. 9 on a lattice. Consider a SU(2N) antiferromagnet on a diamond lattice, where there are two flavors on each site, and the spin of each flavor on every site carries a self-conjugate representation of SU(2N) (Fig. 1c). The spin operator can be represented using slave fermion $f_{i,A,\alpha,a}$:

$$S^{\alpha}_{i,\beta,a} = \sum_{A=1}^{2} \frac{1}{2N} f^\dagger_{i,A,a,\alpha} f_{i,A,a,\beta} - \frac{1}{2N} \delta_{\alpha\beta}. \tag{17}$$

Here $A = 1, 2$ is a color index, $\alpha, \beta = 1, \cdots 2N$ are the SU(2N) indices, $a = 1, 2$ denotes the two flavors. In order to match the spin Hilbert space and fermion Hilbert space, again we need to impose two constraints $\sum_{A,\alpha} f^\dagger_{i,A,a,\alpha} f_{i,A,a,\alpha} = 2N$, and $f^\dagger_{i,A} \rho^\mu_{AB} f_{i,B} = 0$, whose effects can be effectively described by a dynamical compact U(1) gauge field, and a SU(2) gauge field that couples to the color space of $f_{i,A,a,\alpha}$ [28, 29].

Instead of constructing the spin Hamiltonian, let us just consider a spin state, where the slave fermion $f_{i,A,a,\alpha}$ fills a similar mean field band structure as the Fu-Kane-Mele (FKM) model on the diamond lattice, and the flavor index $a = 1, 2$ plays the role of spin in the FKM model [5]:

$$\begin{aligned} H_0 \ = \ & \sum_{<i,j>,A,a,\alpha} -t_{ij} f^\dagger_{i,A,a,\alpha} f_{j,A,a,\alpha} \\ + \ & i\lambda \sum_{\ll i,j \gg} f^\dagger_{i,A,a,\alpha} \vec{\sigma}_{ab} \cdot (\vec{d}^1_{ij} \times \vec{d}^2_{ij}) f_{j,A,b,\alpha}. \end{aligned} \tag{18}$$

The spin wave function is obtained after projecting the slave fermion mean field wave function to satisfy the gauge constraints:

$$|G_{\text{spin}}\rangle = \prod_i \mathrm{P}(\hat{n}_i = 2N) \otimes \mathrm{P}(\hat{\rho}^\mu_i = 0) |f_{i,A,a,\alpha}\rangle. \tag{19}$$

In Eq. 18, when t_{ij} is uniform and isotropic, there are three independent 3d Dirac fermions in the Brillouin zone; two of the three Dirac fermions can be trivially gapped out with an anisotropic t_{ij}, now we are left with one Dirac fermion at $\vec{Q} = (0, 0, \pi)$, whose low energy Hamiltonian reads

$$H = \sum_{\alpha=1}^{2N} \sum_{A} \psi^\dagger_{A,\alpha} \vec{\Gamma} \cdot \vec{q} \psi_{A,\alpha},$$
$$\Gamma_1 = \tau^z \sigma^x, \quad \Gamma^2 = \tau^z \sigma^y, \quad \Gamma_3 = \tau^y. \quad (20)$$

Now we turn on a topological mass gap H_4 to the mean field Dirac Hamiltonian:

$$H_4 = \sum_{\alpha,A,B} m \psi^\dagger_{A,\alpha} \Gamma_4 \rho^z_{AB} \psi_{B,\alpha}, \quad \Gamma_4 = \tau^x. \quad (21)$$

With this mass, the band structure of the slave fermions with color index $A = 1$ is a topological insulator, thus $A = 1$ fermions have massless $2d$ Dirac fermion edge states; the slave fermion with color index $A = 2$ is a trivial band insulator without edge states. H_4 also breaks the SU(2) gauge field down to another U(1) gauge field generated by ρ^z, *i.e.* with nonzero H_4, the slave fermion f is coupled to two different compact U(1) gauge fields at low energy.

Notice that in the current situation we did not particularly assume the time-reversal symmetry in our model, thus one may expect the system to develop another mass gap $H_5 = m_5 \psi^\dagger \Gamma_5 \psi$, where $\Gamma_5 = \tau^z \sigma^z$. Indeed, in the original FKM model, this is precisely the mass gap that breaks the time-reversal and drives the system into a trivial insulator. However, the physical meaning of this mass gap is a slave fermion density modulation between two flavors and two sublattices, in the FKM model it is a spin density wave. Since our original spin model requires a constant slave fermion number on each flavor, then after gauge projection this mass gap does not correspond to any physical order parameter in our lattice spin model.

To build the connection with the NLSM Eq. 9, we couple the slave fermions with the antiferromagnetic Néel order parameter $\mathcal{P}$:

$$H_6 = m_6 \psi^\dagger_\alpha \Gamma_5 \psi_\beta \mathcal{P}_{\alpha\beta}. \quad (22)$$

Both H_4 and H_6 are mass gaps of the Dirac fermion. Now the entire Lagrangian of the Dirac fermion can be written concisely as :

$$\mathcal{L} = \bar{\psi} \gamma_\mu \partial_\mu \psi + \bar{\psi} \mathcal{U} \psi,$$
$$\mathcal{U} = \cos(\theta) \Gamma_4 \rho^z + \sin(\theta) \Gamma_5 \mathcal{P}. \quad (23)$$

$\mathcal{U}$ is a unitary matrix. After integrating out the Dirac fermions, a 3+1d WZW term for unitary matrix U is generated [23]:

$$\mathrm{WZW}(\mathcal{U}) = \int_0^1 du \int d^3x d\tau \frac{2\pi}{480\pi^3} (\mathcal{U}^\dagger d\mathcal{U})^5. \quad (24)$$

Once we assume there is a nonzero background H_4, the $\Theta-$term in Eq. 9 can be derived by directly plugging $\mathcal{U} = \cos(\theta) \Gamma_4 \rho^z + \sin(\theta) \Gamma_5 \mathcal{P}$ in this WZW term Eq. 24.

With a nonzero H_4 in the bulk, the boundary of the system is described by $2N$ two-dimensional Dirac fermions with a SU($2N$) symmetry, and these Dirac fermions are coupled to two U(1) gauge fields. The SU(N) PCM at the boundary (Eq. 10) can also be directly derived using the boundary Dirac fermions, once we break the SU($2N$) symmetry to SU(N)×SU(N) $\rtimes Z_2$. With the Z_2 symmetry, the derived PCM model has precisely $\Theta' = \pi$.

If the Z_2 symmetry at the boundary is further broken, a mass term can be turned on at the boundary: $H_7 = m_7 \bar{\psi} \Omega \psi$. Just like the ordinary 3d topological insulator, this mass term m_7 drives the edge states into a quantum Hall state with Hall conductivity ±1/2 for the SU(N)×SU(N) charges respectively. With absence of topological degeneracy, a fractional Hall conductivity can only occur at the boundary of a 3d system. A domain wall of m_7 is an analogue of domain wall of Θ' in Fig. 1*b*, and using the slave fermions it is straightforward to show that there are nonchiral gapless states localized at the domain wall of m_7.

So far we have ignored the dynamical gauge fields, which in 3+1 dimensional space-time can have a gapless photon phase. In order to make sure the bulk is a fully gapped SPT, we need to drive the system into the confined phase of the U(1) gauge fields. Confinement of U(1) gauge field is driven by condensation of the magnetic monopoles. In the free electron case, a monopole in a topological band insulator will carry gauge charge due to the topological $\theta-$term in the electromagnetic response function [30]. The $\theta-$term leads to "oblique confinement" after the monopoles condense [31, 32]. But in a system where charges are strongly interacting, the quantum number of the lightest monopole, as well as the nature of its confinement transition is not obvious, and I will leave this to future studies. Condensate of bound state between monopole and gauge charges in strongly interacting system is currently under study by other authors [33], and here it is assumed that an ordinary confined phase is still possible by condensing appropriate bound state of monopole and gauge charges. The nontrivial edge physics of the 3d SPT will survive the confinement, because for example the 1+1d CFT at the domain wall in Fig. 1*b* cannot be gapped out without backscattering between left and right moving modes, *i.e.* the domain wall CFT is always stable unless the SU(N)$_L$×SU(N)$_R$ symmetry is spontaneously broken down to its diagonal SU(N) subgroup.

Just like the 1d Haldane phase, the state described above is a 3d SPT *only* if the color singlet constraint $f^\dagger_i \rho^\mu f_i = 0$ is strictly imposed on every site. By contrast, if this constraint is softened, namely the representation on every site is no longer the one in Fig. 1*c*, another mass term can be added to the mean field band structure of the slave fermion: $H_8 = m_8 \psi^\dagger \Gamma_5 \rho^z \psi$, which will completely destroy the bulk SPT, and gap out the edge states without degeneracy. m_8 corresponds to a "color density wave" on the lattice, which is not allowed with the on-site color singlet constraint.

Our construction only applies to the self-conjugate representation in Fig. 1*c*, which is invariant under the center Z_{2N} of group SU($2N$), while the fundamental representation is not invariant under the center Z_{2N}. Thus the 3d SPT state constructed in this work *cannot* be classified using the cohomology of SU($2N$) group, it might be classified using the cohomology of SU($2N$)/Z_{2N}.

3.3 Edge states

Let us take $N = 2$ as an example, and reinvestigate the domain wall configuration in Fig. 1*b*. Let us couple the $\mathrm{SU}(2)_L$ charges to a U(1) gauge field $A_\mu \sigma^z$, which is a spin gauge field that couples to σ^z only. Based on the gapless domain wall states, one can show that if a $2\pi-$flux of A_μ is inserted at the origin $\vec{x} = 0$, this flux will acquire gauge charge 2 [11, 34].

As a comparison, let us make a similar domain wall in a pure 2d system described by the same SU(2) PCM Eq. 10 with $\mathrm{SU}(2)_L \times \mathrm{SU}(2)_R \rtimes Z_2$ symmetry, and inside the domain wall $\Theta' = 2\pi$, where the $\mathrm{SU}(2)_L$ and $\mathrm{SU}(2)_R$ charges have opposite Hall conductivities. Since there is a Z_2 transformation connecting the two different sides of the domain wall, and the Z_2 symmetry exchanges $\mathrm{SU}(2)_L$ and $\mathrm{SU}(2)_R$, thus the systems inside and outside the domain wall should have *opposite* Hall conductivities of A_μ. Since for a 2d bosonic system without fractionalization and topological degeneracy, the Hall conductivity can only be an even integer [11, 12, 34], then inserting a $2\pi-$flux of A_μ inside the domain wall will at least accumulate 4 gauge charges. This proves that with the $\mathrm{SU}(2)_L \times \mathrm{SU}(2)_R \rtimes Z_2$ symmetry, the domain wall states at the edge of the 3d SPT described in Eq. 9 cannot be realized in a 2d system.

Without the Z_2 symmetry that connects the two sides of the domain wall, the argument above would fail. For example, the 2+1 dimensional PCMs discussed in Ref. [34] have no such Z_2 symmetry that connects $\Theta' = 2\pi$ and $\Theta' = 0$.

The same argument can be generalized to the case with $N > 2$. Now we conclude that Eq. 9 describes a 3*d* SPT protected (at least) by the subgroup $\mathrm{SU}(N) \times \mathrm{SU}(N) \rtimes Z_2$ of the SU($2N$) spin symmetry. Since $\mathrm{SU}(N) \times \mathrm{SU}(N) \rtimes Z_2$ is a subgroup of SU($2N$), the original SU($2N$) invariant model Eq. 9 should also describe a 3*d* SPT.

4. DISCUSSION AND SUMMARY

In this work, we studied one class of 3d symmetry protected topological phase, whose stability requires spin symmetry SU($2N$) or its subgroup $\mathrm{SU}(N) \times \mathrm{SU}(N) \rtimes Z_2$, but does not require other discrete symmetries. For large enough N, homotopy groups $\pi_{2d}[\frac{\mathrm{U}(2N)}{\mathrm{U}(N)\times\mathrm{U}(N)}]$ and $\pi_{2d-1}[\mathrm{SU}(N)]$ are always $\mathbb{Z}$ for $d \geq 2$, thus our formalism and results can be generalized to any odd spatial dimension. A $\Theta-$term can be defined for Néel order parameter $\mathcal{P} \in \frac{\mathrm{U}(2N)}{\mathrm{U}(N)\times\mathrm{U}(N)}$ in any odd spatial dimension. After breaking the SU($2N$) symmetry to $\mathrm{SU}(N) \times \mathrm{SU}(N) \rtimes Z_2$ symmetry, this bulk $\Theta-$term always reduces to a boundary $\Theta'-$term with $\Theta' = \pi$ in the SU(N) PCM at the boundary.

This work is supported by the Alfred P. Sloan Foundation, Hellman Family Foundation, and NSF Grant No. DMR-1151208. While preparing for this work, the author became aware of another independent work on 3d SPT [13].

[1] X. Chen, Z.-C. Gu, Z.-X. Liu, and X.-G. Wen, arXiv:1106.4772 (2011).

[2] C. L. Kane and E. J. Mele, Physical Review Letter **95**, 226801 (2005).

[3] C. L. Kane and E. J. Mele, Physical Review Letter **95**, 146802 (2005).

[4] B. A. Bernevig, T. L. Hughes, and S.-C. Zhang, Science **314**, 1757 (2006).

[5] L. Fu, C. L. Kane, and E. J. Mele, Phys. Rev. Lett. **98**, 106803 (2008).

[6] J. E. Moore and L. Balents, Physical Review B **75**, 121306(R) (2007).

[7] R. Roy, Physical Review B **79**, 195322 (2009).

[8] C. Xu and J. E. Moore, Phys. Rev. B **73**, 045322 (2006).

[9] B. A. B. Congjun Wu and S.-C. Zhang, Phys. Rev. Lett. **96**, 106401 (2006).

[10] C. Xu, Phys. Rev. B **81**, 020411 (2010).

[11] M. Levin and T. Senthil, arXiv:1206.1604 (2012).

[12] Y.-M. Lu and A. Vishwanath, arXiv:1205.3156 (2012).

[13] A. Vishwanath and T. Senthil (arXiv:1209.3058).

[14] F. D. M. Haldane, Phys. Lett. A **93**, 464 (1983).

[15] F. D. M. Haldane, Phys. Rev. Lett. **50**, 1153 (1983).

[16] T.-K. Ng, Phys. Rev. B **50**, 555 (1994).

[17] H. Levine, S. B. Libby, and A. M. M. Pruisken, Phys. Rev. Lett. **51**, 1915 (1983).

[18] H. Levine, S. B. Libby, and A. M. M. Pruisken, Nucl. Phys. B **240**, 30, 49, 71 (1984).

[19] H. Levine, S. B. Libby, and A. M. M. Pruisken, Nucl. Phys. B **240**, 49 (1984).

[20] H. Levine, S. B. Libby, and A. M. M. Pruisken, Nucl. Phys. B **240**, 71 (1984).

[21] C. Xu and A. W. W. Ludwig, arXiv:1112.5303 (2011).

[22] C. Xu, F. Wang, Y. Qi, L. Balents, and M. P. A. Fisher, Phys. Rev. Lett. **108**, 087204 (2012).

[23] A. G. Abanov and P. B. Wiegmann, Nucl. Phys. B **570**, 685 (2000).

[24] N. Read and S. Sachdev, Phys. Rev. B **42**, 4568 (1990).

[25] N. Read and S. Sachdev, Nucl. Phys. B **316**, 609 (1989).

[26] E. Witten, Commun. Math. Phys. **92**, 455 (1984).

[27] V. G. Knizhnik and A. B. Zamolodchikov, Nucl. Phys. B **247**, 83 (1984).

[28] C. Xu, Phys. Rev. B **81**, 144431 (2010).

[29] M. Hermele, V. Gurarie, and A. M. Rey, Phys. Rev. B **103**, 135301 (2009).

[30] X.-L. Qi, T. L. Hughes, and S.-C. Zhang, Phys. Rev. B **78**, 195424 (2008).

[31] G. 't Hooft, Nucl. Phys. B **190**, 455 (1981).

[32] J. L. Cardy and E. Rabinovici, Nucl. Phys. B **205**, 1 (1982).

[33] M. P. A. Fisher (private communications).

[34] Z.-X. Liu and X.-G. Wen, arXiv:1205.7024 (2012).